

Module 4. Fluid requirements for football

Unit 4.1 An introduction to fluid and electrolytes for football

Every day, water is lost from the players' body, predominantly in the form of sweat, urine, and respiratory losses, whilst water is gained by the ingestion of foods and fluid in the players' diet. The kidneys regulate water balance by adjusting urine output. Although the total body water content is remarkably constant, water turnover can be very high in some conditions, such as when players train or compete in hot environments. In order to maintain fluid balance, water intake should be adjusted to offset fluid losses on a daily basis (Armstrong *et al.*, 1998; Chevront & Kenefick, 2014; Kavouras *et al.*, 2012). In this unit, we discuss fluid regulation in the player's body before football-specific recommendations in unit 2.


Player's body water

Water has numerous functions in the player's body. Water acts as a solvent for organic and inorganic materials, it provides a medium for biochemical reactions, and it transports solutes throughout the body among various tissues, supplying nutrients and removing waste. The water content of the player's body will range from 45-75%, and it will depend on body composition, age, and gender. The variation of body water content due to body composition is because of muscle (fat-free mass). Muscle has a much higher water content (approximately, 70%-80%) in comparison to fat tissue (approximately, 10%).

The player's total body water can be divided into two compartments: the intracellular water and the extracellular water. The intracellular compartment accounts for approximately 55%-65% of total body water, while the extracellular compartment accounts for the remaining 35%-45% (Chevront & Kenefick, 2014). The extracellular space can be further divided into the interstitial and intravascular fluid (~7.5% of total body water) compartments. Because water is a major component of vascular volume (blood volume is about 35%-45% red blood cells and 55%-65% water [plasma]), hydration also plays a critical role in cardiovascular function and body temperature regulation (Baker & Jeukendrup, 2014). Figure 1 illustrates the body fluid distribution in a 70-kg player.



Figure 1: Body fluid compartments that comprise 42 L of total body water in a 70-kg player and sources of fluid gain or loss



Source: own source.

DID YOU KNOW?

Interstitial fluid is a thin layer of fluid that surrounds the body's cells.

Electrolytes

An electrolyte is a substance that produces a positive or negative charge when dissolved in water. Sodium, potassium, chloride, calcium, magnesium, and phosphate are all examples of electrolytes. Sodium (a cation) and its associated anions (chloride and bicarbonate) comprise the most osmotically active components of the extracellular fluid. Consequently, sodium balance plays a key role in determining the volume of the extracellular fluid compartment and the passive water movement. Sodium achieves this by governing the osmotic gradients between the intracellular and extracellular water spaces.


The most abundant cations in the intracellular water are potassium and magnesium. The imbalance of sodium and potassium across the fluid compartments is maintained by the sodium-potassium pump. All cell functions and electrical communications throughout the player's body are reliant upon maintaining the distribution of electrolytes between the intra- and extracellular fluid (Jain, 2015).

DID YOU KNOW?



Electrolytes can carry a positive or negative charge. An electrolyte which is positively charged is called a cation, and an electrolyte which is negatively charged is called an anion.

Figure 2: Daily fluctuation of a player's body water and the associated terminology



Source: own source.

Temperature regulation

Muscle contraction is relatively inefficient, with approximately 70-75% of energy converted to heat. Therefore, intense exercise is associated with a high level of metabolic heat production (Shirreffs *et al.*, 2005). For every litre of oxygen consumed during exercise, approximately 16 kJ (4 kcal) of heat is produced and only 4 kJ (1 kcal) is actually used to perform mechanical work. During football-specific exercise, some of this heat is stored by the player's body and results in an elevation in core temperature (38-40 °C). If this heat is not dissipated, players would soon overheat and cease to exercise (Ekblom, 1986).

Rises in the body temperature are sensed by central and skin thermoreceptors, and this sensory information is processed by the hypothalamus to trigger appropriate effector responses (Gleeson, 1998).


There are several mechanisms to dissipate heat and to maintain body core temperature in a relatively narrow range both at rest (36-38°C) and during exercise (38-40°C). Although radiation and convection (skin blood flow) contribute, it is the evaporation of sweat that is the most effective mechanism to dissipate heat from the body during exercise (Armstrong *et al.*, 1997; Chevront & Kenefick, 2014). Sweat must evaporate from the body surface in order to exert a cooling effect, with the evaporation of 1 L of sweat from the skin, removing 2.4 MJ of heat from the body (Wenger, 1972). Although sweating is a very effective way to dissipate heat, it may cause hypohydration if sweat losses are not replenished (Chevront & Kenefick, 2014; Chevront *et al.*, 2013). Hypohydration has the

potential to impact on performance, which may have implications for day-to-day training and match performance.

DID YOU KNOW?

The hypothalamus is a small region, located at the base of the brain. The hypothalamus is pivotal in regulating the players' body temperature but also functions to regulate various other physiological functions, including releasing hormones, appetite and emotional responses.

Figure 3: A schematic summarising the process of heat loss and sweat evaporation from skin surface to create cooling



Source: own source.

Dehydration and exercise performance

As the player's body becomes progressively hypohydrated, a reduction in skin blood flow and sweat rate may occur, impairing the ability to regulate body temperature. Several studies have shown that mild hypohydration, equivalent to the loss of 2% body mass (BM)

can be sufficient to impair exercise performance (ACSM, 2016). In addition, it is often reported that greater losses in BM will result in greater reductions in performance (Baker *et al.*, 2007; Sawka *et al.*, 2012), especially in hot conditions. Although the detrimental effects of hypohydration on performance are most pronounced in hot conditions, such effects have also been observed in cool conditions (Cheuvront *et al.*, 2005). Others have argued that performance may not be affected by mild hypohydration (Goulet, 2011). It is highly likely that there are individual differences among players in their ability to tolerate differing levels of hypohydration. Therefore, there may be discussion about the exact level of hypohydration that is acceptable in football players. Sweat losses do not often exceed 2% of pre-exercise body mass during training. However, during matches, sweat losses may be greater, especially when matches are played in the heat. Based on the available literature, 2-3% hypohydration seems a practical and safe allowance for football players. It is important to note that some players may be at greater 'risk' of significant hypohydration >2-3% BM losses, which will be discussed in unit 2.

KEY POINT

Dehydration is the process of losing body water.

Hypohydration is a deficit of body water that is caused by acute or chronic dehydration.

Both decreases in maximal aerobic power and decreases in endurance capacity have been reported with hypohydration in temperate conditions (McConnell *et al.*, 1997), although not all studies found such an effect (Maughan *et al.*, 1989; Robinson *et al.*, 1995). There are several reasons why hypohydration results in decreased exercise performance. The most dramatic consequence of dehydration-induced (5% pre-exercise BM loss) hyperthermia during exercise in extreme conditions (35 °C) is a 25-30% reduction in stroke volume that is not generally met with a proportional increase in heart rate. This results in a decline in cardiac output and in arterial blood pressure (Gonzalez-Alonso *et al.*, 1995). In addition, during exercise in the heat, the dilation of the skin blood vessels reduces the proportion of the cardiac output that is devoted to perfusion of the working muscles.

Becoming significantly hypohydrated will also impair the ability of the player's body to lose heat. Both sweat rate and skin blood flow are lower at the same core temperature when hypohydrated in comparison to when the individual is in a hydrated state. This means that body temperature rises faster during exercise when the body is hypohydrated, and this is commonly accompanied by a higher heart rate during exercise (Coyle & Gonzalez-Alonso, 2001; Montain & Coyle, 1992). Finally, a larger rise in core temperature during exercise in the hypohydrated state is associated with an increased rate of muscle glycogen breakdown (Hargreaves *et al.*, 1996; Logan-Sprenger *et al.*, 2015). Depletion of glycogen stores could also result in premature fatigue and reductions in



high-intensity running towards the end of a football match (Bendiksen *et al.*, 2012). In addition to the effects of hypohydration on endurance performance, there are also reported negative effects on perception of effort, coordination, and cognitive functioning (Baker *et al.*, 2007; Logan-Sprenger *et al.*, 2015). This is likely to negatively impact a player's skill and decision-making process.

It was previously thought that muscle strength and anaerobic performance were less likely to be affected by hypohydration (Ali & Williams, 2013; Chevront & Kenefick, 2014). However, a meta-analysis revealed that hypohydration, or factors associated with dehydration, are likely to be associated with practically important decrements in muscle endurance, strength, and anaerobic power and capacity (Savoie *et al.*, 2015).

It would be intuitive to consider whether reductions in body mass through dehydration during football exercise could be advantageous to sprint speed and vertical jump height (Viitasalo *et al.*, 1987). However, in one study, a diuretic-induced reduction in body mass of 2.5% had no effect on sprint or power performance (Watson *et al.*, 2005). Likewise, there was no correlation between the reduction in body mass and vertical jump height (Watson *et al.*, 2005). These results suggest that hypohydration provides no advantage for weight-bearing activities required for football. Some have argued that 'faster' runners lose more body mass during exercise (Knechtle *et al.*, 2012; Knechtle *et al.*, 2012; Rust *et al.*, 2012). Whilst this may be true, it is important to note that faster runners will have higher sweat rates. Furthermore, these are simply correlation studies that do not show causality, and it is even possible that individuals who can tolerate larger fluid losses are more competitive.

Football requires not only aerobic performance, but also a sustained level of technical and skill proficiency throughout exercise (Harper *et al.*, 2014; Russell & Kingsley, 2014). Owen *et al.* (2013) examined the effect of dehydration on football skills (i.e., passing and shooting) and intermittent high-intensity running performance after the 90-minute Loughborough Shuttle Running Test (LIST) protocol in a temperate environment (19 °C, 59% relative humidity). Despite differences in fluid intake (no fluid, *ad libitum*, and prescribed volume) and hypohydration (2.5%, 1.1%, and 0.3%, respectively), football skills and intermittent high-intensity exercise performance were similar after the LIST (Owen *et al.*, 2013). These results were in contrast to McGregor and colleagues (McGregor *et al.*, 1999), who were the first to test the effects of dehydration on football-specific performance. In this study, ratings of perceived exertion were higher toward the end of the 90-minute LIST (13-20 °C, 57% relative humidity [RH]) when no fluid was given to the players (resulting in 2.5% hypohydration), compared to when fluid was provided (resulting in 1.4% hypohydration). Likewise, 2.5% hypohydration slowed sprint time at the end of the LIST in comparison to 1.4% hypohydration. This study also showed that football-specific skill performance (i.e. dribbling skill) decreased by 5% from pre- to post-LIST with 2.5% hypohydration, but it was maintained when hypohydration was limited to



1.4% pre-exercise body mass. However, 2.5% hypohydration had no impact on football players' mental concentration test scores at the end of the LIST (McGregor *et al.*, 1999).

Morris *et al.* (2005) later reported that prolonged, intermittent, high-intensity shuttle running in the heat (33 °C, 28% RH) resulted in earlier onset of exhaustion in comparison to exercise in a moderate environment (17 °C, 63% RH). Interestingly, while muscle glycogen utilization was elevated by heat stress, low muscle glycogen concentration was not reported to be the cause of the earlier exhaustion. Instead, the onset of exhaustion was associated with hyperthermia (Morris *et al.*, 2005). In support of these findings, a study which asked football players to complete a treadmill-based football protocol at 18 °C and 30 °C revealed total distance covered and sprint distances to be significantly reduced when exercise was performed in the hot environment (Aldous *et al.*, 2015). Specifically, the reduction in sprint distance in hot conditions was accompanied by higher heart rate and an elevated core temperature (~0.4 °C) (Aldous *et al.*, 2015). Therefore, maintaining the ability to thermoregulate and reduce the rate of rise in core temperature is of benefit to football performance. The importance of thermoregulation for football players is further exacerbated in matches held in warm environments that exceed 90 minutes, i.e. in extra time (Harper *et al.*, 2014).

Interestingly, it is highly likely that individual players may be more or less sensitive to hypohydration. The magnitude of exercise-heat stress 'fatigue/response' has high inter-individual variation due to training and acclimation status, which are influenced by genetic/phenotypic variations of favourable traits associated with innate thermal tolerance and its acquirement (Horowitz, 2014; Nybo *et al.*, 2014). Higher muscle temperatures within the quadriceps and elevated core temperatures have been reported when playing football in the heat, compared to temperate environments (Mohr & Krstrup, 2013). Well-trained players will be able to tolerate higher core temperatures than less-trained individuals (up to 40.3 °C). The improved heat tolerance coincides with prolonged time to exhaustion during constant paced exercise (Cheung, 2010; Sawka *et al.*, 2011).

One theory is that the degree of hypohydration which can be tolerated by the player may be a result of the degree of hypohydration which the player has become accustomed to or routinely experiences. Research in cycling demonstrated that as little as four familiarisation sessions, designed to habituate the individual with the dehydration protocol, attenuated the performance decrement observed when 2.4% hypohydrated (Fleming & James, 2014). Nevertheless, it is important to note that performance following habituation was still lower compared to when exercise was completed hydrated. Interestingly, there were no physiological mechanisms to explain the results (i.e., familiarisation did not attenuate cardiovascular strain), so the advantage was likely a result of improved perceptual/psychological response (Rating of perceived Exertion [RPE]). Although this study was performed in cycling and football-specific research is yet



to be completed, players and coaches are advised to not allow hypohydration to become the 'norm' and become accustomed to consistent sub-optimal performance. That being said, this study also highlights the importance of routinely exposing players to the level of hypohydration that simulates competitive demands (unit 2).

Determinants of fluid intake

The net body water balance of the player (fluid losses = fluid gains) is regulated remarkably well day-to-day, as a result of their thirst and hunger combined with the access to food and beverages. However, when fluid losses are greater than fluid intake, it results in hypohydration. Because a player's body water has a normal daily fluctuation, hypohydration may be defined as a body water deficit greater than normal daily fluctuation (Cheuvront & Kenefick, 2014). Hypohydration may also be defined as a negative change in body water which exceeds 2 percent of the normal body mass variability (Cheuvront *et al.*, 2011).

When a football player is at rest, the approximate threshold level of hypohydration where compensatory fluid regulatory actions (fluid conservation at the kidney) and stimulus for fluid acquisition (thirst) occurs is approximately >2% of their body mass (Shirreffs *et al.*, 2004). These compensatory actions are triggered by elevations in blood plasma osmolality and, to a lesser degree, a reduction in plasma volume. During football exercise, particularly in the heat, plasma volume decreases because it provides the fluid for sweat, and, as a result, plasma osmolality increases because sweat has a low sodium content relative to blood plasma (Kenefick, 2018).

It is important to remember that, when total body water losses occur during football exercise due to thermoregulatory sweating, these losses are shared by all fluid compartments (Kenefick, 2018). An increase in plasma and extracellular osmolality will pull fluid from the intracellular space so that all compartments are in non-osmotic equilibrium. A ~2% increase in plasma concentration is commonly referenced as an osmotic threshold for compensatory water retention at the kidneys and water acquisition (thirst), which is also equivalent of ~2% BM loss. These losses would be the equivalent of 1.4 L of fluid for a 70-kg player.

Metabolic water production can also contribute to water gain during exercise. Water is produced as a result of aerobic metabolism. The rate of water formation is primarily dependent upon energy expenditure and, to a lesser extent, upon the type of substrate. The water produced through metabolism has been estimated to be approximately 250 to 350 mL per day for sedentary persons and up to 500 to 600 mL per day for physically active individuals (Baker & Jeukendrup, 2014). Finally, although only small, it is also important to recognise that sources other than sweat (metabolic mass loss from substrate metabolism and respiratory water losses) may also contribute to body mass losses during exercise (Cheuvront & Montain, 2017).



Table 1: The water content range for selected foods

Percentage	Food item
100%	Water
90% - 99%	Fat-free milk, cantaloupe, strawberries, watermelon, lettuce, cabbage, celery, spinach, pickles, squash (cooked)
80% - 89%	Fruit juice, yogurt, apples, grapes, oranges, carrots, broccoli, (cooked), pears, pineapple
70% - 79%	Bananas, avocados, cottage cheese, ricotta cheese, potato (baked) corn (cooked) shrimp
60% - 69%	Pasta, legumes, salmon, ice cream, chicken breast
50% - 59%	Ground beef, hot dogs, feta cheese, tenderloin steak (cooked)
40% - 49%	Pizza
30% - 39%	Cheddar cheese, bagels, bread
20% - 29%	Pepperoni sausage, cake, biscuits
10% - 19%	Butter, margarine, raisins
1% - 9%	Walnuts, peanuts (dry roasted), chocolate chip cookies, crackers, cereals, pretzels, taco shells, peanut butter
0%	Oils, sugars

Source: Popkin *et al.*, 2010, p. 25.

Monitoring hydration status


Assuming the player is in energy balance, daily hydration status may be estimated by tracking early morning body weight (measured upon waking and after voiding), since acute changes in body weight generally reflect shifts in body water (Thomas *et al.*, 2016). Urine analysis may also be a useful indicator of a player's hydration status. Urinary-specific gravity and urine osmolality may be used to approximate hydration status by measuring the concentration of the solutes in urine (Thomas *et al.*, 2016). Ideally, the urine should be collected from a midstream collection of the first morning urine sample. As a



guide, a urinary-specific gravity of less than 1.020—perhaps ranging to 1.025 to account for individual variability—is generally indicative of euhydration. Urinary osmolality reflects hypohydration when greater than 900 mOsmol/kg, while euhydration is considered as less than 700 mOsmol/kg (Thomas *et al.*, 2016).

Most likely, the easiest method for the player to assess their hydration status is the volume and colour of their urine. If a large volume of urine is produced and the urine is pale yellow in colour, the player is most likely hydrated or drank sufficient fluid with their meals. If the urine is very dark in colour (dark orange/brown), it is likely that they have not drunk enough volumes of fluid following their last bout of football activity or with their last meal. In this case, fluid intake should be adjusted accordingly (unit 2). Practically, placing ‘hydration’ reminders or urine colour charts (Armstrong *et al.*, 2010) above urinals in male toilets or in female cubicles serve as a behavioural ‘nudge’ for players to remember to monitor their hydration status. Likewise, targeted hydration education around pre-season and during periods of fixture congestion are recommended.

Figure 4: Urine colour chart




Source: own creation.

Figure 5: Summary of mechanisms by which hypohydration can impact player performance

Mechanisms

Likely a combination of mechanisms that contribute to reduced endurance performance



Source: adapted from James *et al.*, 2019.

(It will likely be a combination of mechanisms that contribute to reduced football performance).

Summary

- Water accounts for approximately 55-60% of the player's body mass.
- A player's total body water will naturally fluctuate throughout the day.
- The player's total body water can be divided into intracellular water and extracellular water. Sodium is the main electrolyte in extracellular water.
- Fluid is lost during exercise as players sweat to keep the body cool.
- Hypohydration >2% of pre-exercise body mass may negatively impact football-specific performance via a combination of mechanisms.

Unit 4.2 Fluid and electrolytes for football

Every day water is lost from the players' body in the form of sweat, urine, and respiratory losses, whilst water is gained by the ingestion of foods and fluid in the players' diet (unit 1). Hypohydration of >2% body mass deficit has been shown to impair football-specific performance, including intermittent high-intensity sprinting and dribbling skills. It is often observed that football players start a practice or match play already in a hypohydrated state, probably as a result of cumulative dehydration from previous training or matches. Therefore, fluid intake on a daily basis may be as important as fluid intake strategies during competition (we present a summary of both scenarios, i.e. daily habits and exercise). In this unit, we discuss how players can be educated to assess their hydration status, as well as how to determine individual player fluid requirements. Finally, practical advice will be shared on how to deliver the hydration strategy throughout the club.

Player sweat rates

Sweat rates vary greatly among players and are primarily influenced by the intensity of exercise, environmental conditions, and acclimation status (Duffield *et al.*, 2012). During training and matches, sweat rates have been reported to range from 0.5 L/h to 2.5 L/h (Baker *et al.*, 2016; Shirreffs *et al.*, 2005), and are generally lower in female players because of lower body mass and absolute work rates (Kilding *et al.*, 2009). The mechanisms by which sweating-induced hypohydration might impair football performance are not completely understood. However, they may include increased cardiovascular strain (Armstrong *et al.*, 1985), impaired cognitive function (Ganio *et al.*, 2011), increased perception of effort (McGregor *et al.*, 1999), reduced physical function (Mohr & Krstrup, 2013), and reduced technical skills (McGregor *et al.*, 1999). Players are therefore advised to drink sufficient fluids to prevent a deficit of >2% of pre-exercise body mass loss during exercise (Sawka *et al.*, 2007). Conversely, it is important to note that players must avoid gains in body mass (hyperhydration) during exercise.

During training and matches, players should pay attention to ingesting an appropriate quantity of carbohydrate (module 1), whilst simultaneously meeting fluid needs. Both hydration and carbohydrate ingestion will require additional attention in matches where extra-time (2x15 minutes) is played. All match nutrition strategies, including hydration, should be practiced in training and training matches to allow individualised protocols to be developed and to identify potential adverse effects in players. This method also allows players the opportunity to become adapted to the nutrition strategy without impacting on important/competitive match performance.

To highlight the fluid demands experienced by players, if we assume a higher level of total body water (i.e. 50 L) in cool, low-intensity training environment, total body would turn over in approximately 100 playing hours. However, during match play in warm



environments, at a modest sweat rate of 1.5 L/h, total body water would turn over in approximately 33 playing hours. In practical terms, total body water would turn over in just under four weeks for a player completing 2 games a week and completing a 1-hour training session between games. Thus, over a 38-week season, a player's total body water may turn over a conservative ten times.

Method to monitor player fluid balance

It is possible to measure the fluid balance of players across training and matches. By routinely adopting this method, it will help inform the individual players of their fluid needs during exercise and inform the coaches to the level of hypohydration being experienced by the player. Specifically, fluid intake and losses can be recorded and used to calculate the player's sweat rate (Baker, 2016).


It is important to note that body water gain occurs through metabolic water production via the dissociation of water from glycogen and substrate use (unit 1). It is generally accepted that for most exercise (~2 hours or less), evaporative loss of water from the lungs through respiration is negligible, as are changes in metabolic mass, and can be ignored (Cheuvront & Montain, 2017; Maughan *et al.*, 2005). For practical purposes, it is assumed that 1 kg of a player's body mass loss represents approximately 1 L of water loss (Maughan *et al.*, 2007; Maughan *et al.*, 2007). Therefore, sweat loss can be calculated from the change in a player's body mass during exercise following the correction for fluid intake. The method to measure fluid balance during exercise is displayed in figure 4 and summarised below:

1. Collect a pre-exercise (ideally first morning) urine sample from the players. These samples can be used as an indicator of the players' hydration status, either through analysis of urine-specific gravity/osmolality or—more simply—by analysis of volume and colour.
2. Before the first BM measurement, encourage players to void their bladder. Record the players' BM (kg). This can be completed by players wearing minimal clothing (i.e. underwear) or nude if privacy is possible.
3. Allocate each player a drink bottle, which should be weighed before and after exercise. Changes in the bottle weight in grams represents the volume of fluid ingested in mL. Advise the players to only drink from their bottle and to avoid squeezing water onto their faces for cooling purposes.
4. During exercise, record the duration of exercise and the environmental conditions, and collect, measure, and record any urine which is passed by the players.
5. After exercise, ensure the players are dry before the final body mass measurement is recorded (kg). A change in body mass of 1 kg is the equivalent of 1 L of fluid loss.



An example of how to calculate the sweat rate of a player is displayed in figure 4. This method allows for individual sweating response to training and matches to be captured. It also helps the nutritionist identify those players with 'high' sweat rates, and thus at greater risk of significant hypohydration.

Figure 6: Fluid balance method overview



Source: own source.

Figure 7: Player sweat rate calculation and working examples

Sweat loss (L)=

BODY MASS LOSS (kg) + FLUID INTAKE – URINE VOLUME

Sweat rate (L/h)=

SWEAT LOSS (L) / DURATION (HOUR)

Sweat loss (L)=

BODY MASS LOSS (kg) + FLUID INTAKE – URINE VOLUME = 2.0 L

1.5

500 ml / 1000

0 ml / 1000

Sweat rate (L/h)=

SWEAT LOSS (L) / DURATION (HOUR) = ~1.3 L/h

2.0

1.5 (90 mil)

Source: own source.

Sweat analysis

Sweat is hypotonic but contains electrolytes (unit 1). The major electrolytes found in sweat are sodium (Na⁺) and chloride (Cl⁻); collectively, these ions form the salt (sodium chloride). Analysis of sweat composition allows for the amount of sodium chloride lost during exercise to be determined, and differences among individual players to be measured. It is sometimes easy to identify players with high salt losses, as you will be able to see a residue of salt on their kit after training or matches.

The replenishment of sodium chloride after exercise is important for complete and rapid rehydration detailed below (Shirreffs *et al.*, 1996). Full details of the sweat collection and analysis protocol have been described in detail (Baker *et al.*, 2009). In summary, absorbent patches are covered with an adhesive film that holds the patch in place and prevents sweat evaporation. The patches should be placed on the skin surface following washing with deionized water and drying with a sterile gauze swab. Patches remain in place for the duration of exercise and are removed immediately after exercise, or sooner if they become saturated.



There are different methods to analyse the sweat electrolyte composition (Baker *et al.*, 2009). Historically, this analysis has been laboratory based, relatively slow and expensive. Recently, a field method has been validated against gold standard laboratory techniques which allows rapid analysis of sweat composition (Baker *et al.*, 2014). After removal from the skin, patches can be immediately placed inside the barrel of a 5-mL syringe using sterile tweezers. The syringe plunger is then depressed to compress the patch, and the expelled sweat is collected into a sterile tube. Sweat samples can be analysed immediately using a compact wireless analyser that uses ion-selective electrode technology (Horiba B-722) to derive measures of sodium concentration. Sweat electrolyte losses (g) should be normalised, and whole-body losses should be calculated from the sweat electrolyte concentration, the molecular weight of the electrolyte, and the total sweat loss of the player (Baker *et al.*, 2016; Baker *et al.*, 2014).

Scientific innovations in this area included single-use sweat patches that, when paired with a smartphone application, can determine sweat rate and sweat composition after exercise (Baker *et al.*, 2020). Specifically, patches use microfluidic technology to measure sweat rate and a colorimetric assay to determine sweat chloride concentration. As sweat chloride is directly associated with sweat sodium, sodium can be reported. These technological approaches serve as excellent engagement tools with players, providing a quick and easy way to highlight sweat losses, with minimal intervention. As new and real-time sweat rate devices are developed, you are encouraged to ensure that the technology has been validated appropriately.

Fluid intake

To maintain water balance, fluid intake must compensate for the fluid loss that occurs during training, matches, and daily living. Fluid intake is usually dependent on thirst feelings, but thirst (or the lack of thirst) can also be overridden by conscious control (Sawka & Greenleaf, 1992). It is important to note, however, that thirst has been reported to be a poor indicator of fluid requirements or the degree of hypohydration during exercise (McKenna & Thompson, 1998). As discussed, even a mild degree of hypohydration can be sufficient to impair exercise performance, especially during exercise in warm conditions (Cheuvront & Kenefick, 2014). In studies investigating sweat rates in professional football, fluid has been freely available. Those studies which have reported the volume of fluid voluntarily ingested have shown that drinking was highly variable among players in the same teams and was also influenced by the environmental conditions (Maughan & Shirreffs, 2007a, 2007b). Maughan *et al.* (2005) tabulated results from five studies that investigated fluid loss and intake of elite male football players at temperatures ranging from 5-32 °C, and found *ad-libitum* fluid intake ranged from 423-1401 mL (Maughan *et al.*, 2007). Unfortunately, the volume of the fluid intake, which was water and which was a beverage containing carbohydrate, is not commonly reported. This is of interest because the routine ingestion of carbohydrate provided in combination



with fluid, typically via carbohydrate-electrolyte beverages, has been reported to positively influence various aspects of football-specific performance (Rollo, 2014; Russell & Kingsley, 2014) and outcomes associated with team sport performance (Baker *et al.*, 2015; Williams & Rollo, 2015).

DID YOU KNOW?

The term 'ad libitum' simply means self-selected or freely chosen.

Fluid loss

Sweat losses in professional footballers have been reported to vary greatly among players, even in response to the same exercise session (Aragón-Vargas *et al.*, 2009). The two main determinants of the rate at which fluid is lost as a consequence of sweating (sweat rate) are exercise intensity and environmental conditions (Hawley *et al.*, 1994; Maughan *et al.*, 2007). Sweat rate is primarily determined by metabolic heat production, which is directly proportional to absolute exercise intensity. The environmental heat stress is determined by the ambient temperature, relative humidity, wind velocity, and solar radiation. Of these factors, the ambient temperature and relative humidity are considered the most important. A high ambient temperature increases skin blood flow, and high humidity will severely compromise the evaporative loss of sweat. Rather than evaporate, sweat will often drip off the skin in such conditions, diminishing the effectiveness of heat loss via this route.

Football is practised and played in a variety of environmental conditions. In Barcelona, during the same competitive season, players train and compete in environmental conditions ranging from cool in the winter months (wet bulb globe temperature [WBGT], 15 ± 7 °C, $66 \pm 6\%$ RH, January 2015) to very hot in the summer (WBGT $>29 \pm 1$ °C, 52 ± 7 RH, July 2014). The mean sweat rate of professional players training in hot ambient temperatures (32.3 ± 3 °C) has been reported to be 1.46 ± 0.24 L/h (range 1.12-2.09 L/h), and the corresponding mean level of hypohydration was $1.59 \pm 0.61\%$ of pre-exercise body mass (range 0.71-3.16%) (Shirreffs *et al.*, 2005). Although mean sweat losses have been reported to be lower (1.13 ± 0.30 L/h, range 0.71-1.77 L/h) when training in cool temperatures (5.1 ± 0.7 °C), players still experience significant levels of hypohydration: $1.62 \pm 0.55\%$ (range 0.87-2.55%) (Maughan *et al.*, 2005).

Female players will generally have lower sweat rates than their male counterparts due to a smaller body size and less muscle mass contributing to metabolic heat production during exercise (Cheuvront *et al.*, 2002). Because of the lower fluid and electrolyte losses in female players, some studies have suggested that hydration is unlikely to require special consideration in this group (Kilding *et al.*, 2009). However, in our experience, women players' sweat response will also vary significantly among players in response to the same training session (Tarnowski *et al.*, 2022). In addition, it is also a challenge for



female players to drink large volumes of fluid prior to and in recovery from exercise. Thus, more research and insight are needed to optimise the hydration practices of this elite population (course 4).

Case study from FCB

In the 2014-2015 season (in which FC Barcelona men's first team won: La Liga, the Copa del Rey, and the UEFA Champions League), fourteen male first-team players of Fútbol Club (FC) Barcelona (Spanish first division; La Liga) completed a sweat test on four different training occasions (Rollo *et al.*, 2021). Each training session was completed outside on a grass pitch in dry conditions. Environmental temperature and relative humidity were recorded at the beginning and at 15-minute intervals during training. The training was that which was normally conducted by the players at that time of the season. Importantly, testing was achieved with no interference to the players' normal routine. The training intensity was classified by a global positioning satellite system. Low intensity training was classified as players covering 40-46 meters/minute whilst high intensity training was 82-86 meters per minute. In addition, at the cessation of training, players were asked to rate 'how hard' the training session was on a 1-10 rating of perceived exertion (RPE) scale. Ratings of 1-5 were indicative of low intensity, and ratings of 6-10 were indicative of high intensity. Data were collected under the following conditions classified by WBGT: cool low intensity (CL) (14 ± 7 °C, $67 \pm 7\%$ RH); cool high intensity (CH) (14 ± 8 °C, 69 ± 7 RH); hot low intensity (HL) (28 ± 1 °C, 55 ± 9 RH); hot high intensity (HH) (28 ± 2 °C, 55 ± 10 RH). For each trial, there were three drinks breaks per hour of exercise. The duration of each drinks break was 90 ± 30 s. The main variables are listed in full in table 1.

Table 1: Physiological and sweat electrolyte responses to exercise at different intensities and environmental conditions

Intensity	Condition				F	p
	Cool		Hot			
	Low	High	Low	High		
Urine Specific Gravity	1.024 ± 0.005 (1.016-1.032)	1.023 ± 0.004 (1.014-1.028)	1.023 ± 0.006 (1.011-1.033)	1.025 ± 0.005 (1.017-1.034)	(3,52) = 0.35	0.786
Sweat rate (L·h ⁻¹)	0.55 ± 0.20 (0.20-0.85)	0.98 ± 0.21^a (0.67-1.50)	0.81 ± 0.17^a (0.50-1.20)	1.43 ± 0.23^{abc} (1.10-1.81)	(3,52) = 46.37	0.001
Fluid intake (mL·h ⁻¹)	394 ± 160 (184-719)	505 ± 265 (220-1058)	572 ± 214 (308-898)	663 ± 229^a (266-933)	(3,52) = 3.68	0.018
Carbohydrate intake (g·h ⁻¹)	12 ± 9 (0-37)	11 ± 11 (0-32)	15 ± 12 (0-36)	15 ± 14 (0-50)	(3,52) = 0.57	0.637
Sweat [Na ⁺] (mmol·L ⁻¹)	35 ± 9 (21-54)	38 ± 8 (26-48)	34 ± 7 (24-45)	38 ± 8 (22-53)	(3,52) = 0.85	0.475
Sweat Na ⁺ loss (mmol·h ⁻¹)	19 ± 9 (7-36)	38 ± 13^a (19-68)	28 ± 9^a (13-52)	54 ± 15^{abc} (33-81)	(3,52) = 22.98	0.001
NaCl loss (g·h ⁻¹)	1.1 ± 0.5 (0.4-2.1)	2.2 ± 0.8^a (1.1-3.9)	1.6 ± 0.5^a (0.8-3.0)	3.2 ± 0.9^{abc} (1.9-4.8)	(3,52) = 22.98	0.001

Source: Rollo *et al.*, 2021, p. 5.

Table 1 description:

“The mean \pm s and (range) of variables recorded before and during each training session in response to exercise at low and high exercise intensities in cool and hot conditions. Predicted whole-body sweat [Na⁺], predicted whole-body sweat Na⁺ losses, and assumed sodium chloride losses during exercise. The loss of sodium chloride is calculated on the assumption that all Na⁺ loss is sodium chloride [27]. a = significantly different from Cool Low, b = significantly different from Cool High, c = significantly different from Hot Low” (Rollo *et al.*, 2021, p. 5).

From a practical perspective, following each testing session players received personalised feedback, which was refined as more tests were completed. During each training session, players drank sufficient volumes of fluid to prevent body mass losses in excess of 2% pre-exercise body mass values. The results in table 2 demonstrate the variability in sweat rate and sweat sodium concentrations in this population of elite professional players. Interestingly, the results of the tests showed that sweat sodium concentration did not change between different sessions. Perhaps, sweat sodium may only need to be completed on a single occasion over the season. However, further research is required to substantiate this observation.

Despite these observations for training, greater challenges to fluid balance occur during matches, especially in hot conditions, as sweat rates (water losses) increase dramatically (Tarnowski *et al.*, 2022). High sweat rates combined with limited opportunities to drink during competitive match play often result in players experiencing significant hypohydration during matches (>2% BM)(unpublished match day observations).

Fluid intake strategies

Fluid intake during training and matches can help maintain plasma volume and prevent the adverse effects of dehydration on endurance and technical performance. When there is only little time in between two training sessions, rapid rehydration is crucial, and drinking regimes need to be employed to optimise fluid delivery. As such, strategies for fluid replacement before, during, and after football exercise will be discussed in the following sections.

Fluid intake before training and matches

The aim for the player should be to begin training and games hydrated. Generally, there is sufficient time, i.e. >8 hours between training sessions or matches, for the player to consume sufficient volumes of fluid with meals to achieve this. However, during matches, the player may incur in significant fluid losses, and thus more structured hydration strategies may be required. An effective pre-exercise hydration strategy will help ensure



that any previous fluid deficits that the player has experienced from exercise or travel be rectified before re-engaging in football-specific activity.

As a guide, to ensure adequate hydration prior to exercise, players are advised to drink a volume of fluid specific for their body mass. Specifically, at least four hours before exercise, players should gradually drink approximately 5-7 mL of fluid per kg body mass. On match days, this can be completed with their final meal before the match, whereas on training days, these volumes should be ingested with or before breakfast. Consuming fluid with meals with small amounts of sodium or sodium-containing foods will help stimulate thirst and the retention of the ingested fluid (Maughan & Leiper, 1995). Drinking approximately four hours before exercise allows sufficient time for urine output to be expelled before players are expected to perform. In addition, it avoids the situation of players drinking too much fluid immediately before exercise, which increases the risk of the player feeling bloated and experiencing gastrointestinal complaints.

In a situation where the player does not produce urine, or the urine is of low volume and dark in colour, a further 3-5 mL of fluid per kg body may be advised approximately 2 hours before exercise. It is important to note that fluid intake is, of course, self-regulated by the player. Therefore, adequate education programmes within the club should be in place so that the player can identify signs of hypohydration themselves and modify their drinking behaviour appropriately (unit 1).

Fluid intake during training and matches

In order to avoid significant hypohydration during the match or training, fluids should be ingested. By regularly measuring body mass before and after exercise, it is possible to build a profile of fluid losses in response to specific footballing sessions (i.e. training/games) and conditions (lower or higher temperatures and humidity). Sufficient data collection over time will allow predictions of fluid losses based on exercise intensity and forecasted weather.

Fluid intake is likely to benefit the player when exercise is longer than 30-60 minutes, but there appears little advantage during strenuous exercise of less than 30 minutes in duration. During such high-intensity exercise, gastric emptying is inhibited and the drink is more likely to cause gastrointestinal distress with no performance benefit. Large volumes of fluid are difficult, if not impossible, to ingest, and even one litre may feel quite uncomfortable in the 'untrained' stomach (Murray, 2006). Therefore, in these circumstances, it is not advised nor is it practically possible to match fluid intakes to sweat losses during exercise. Another factor that can make the ingestion of large amounts of fluid difficult is the fact that, in matches, the opportunities for drinking are limited to before the game, half time, and during unscheduled breaks in play (Baker *et al.*, 2016; Gleeson, 1998).



Fluid intake has been found to be a strong predictor of outdoor team sport pre-season training performance. Players that drank a volume of fluid above ~10 mL/kg BM achieved a 4.1% increase in high-speed running distance over training, in comparison to those players who failed to drink equivalent volumes (Badham *et al.*, 2023). However, excessive water intake should be avoided, as this has been linked to hyponatremia (low plasma sodium). Although, to date and to the author's knowledge, there have been no reported cases of hyponatremia in football players, education about the appropriate volumes of fluid to drink (based on sweat rates in combination with few drinking opportunities that exist both in training and on match days) should avoid the problem of over-drinking. Therefore, as a guide, players should aim to minimise fluid losses during exercise of an hour or more to equal to or less than 2% of starting body mass (assuming the player started in an euhydrated state).

When faced with drinking guidelines, some players may be overwhelmed by the volumes of fluid suggested. If a player's current drinking regime is in stark contrast to recommendations, it is advised to work towards fluid intake goals over time. Specifically, do not introduce large changes in one go. Drinking before exercise can be built into a daily routine, gradually increasing the volume. During exercise, it is often difficult to tolerate the volumes of fluid needed to prevent significant hypohydration. However, the volume of fluid that is tolerable can be trained and increased with frequent drinking in training (Lambert *et al.*, 2008). Drink training will accustom the players to the feeling of exercising with fluid in the stomach, whilst helping to optimise the delivery of carbohydrate if sports drinks are chosen (Jeukendrup, 2011). It also gives the player the opportunity to experiment with different volumes and flavourings to determine the volume of fluid intake they can tolerate, and which formulations suit them best.

Type of fluid

Numerous studies have shown that 'regular' water intake during prolonged exercise is effective in improving performance (for review, see Shirreffs & Sawka, 2011). However, fluid intake during exercise also offers the opportunity to provide some fuel in the form of carbohydrate (module 1). The type and composition of drink consumed during exercise should be specific to the exercise session, as well as the goals of each individual player (Impey *et al.*, 2016; Jeukendrup, 2014). Therefore, during match play, where 'performance' is the objective, the addition of carbohydrate to drinks consumed will have an additive and independent effect in comparison to water on exercise performance (Below *et al.*, 1995; Nicholas *et al.*, 1995; Williams & Rollo, 2015).

Beverages for fluid and energy replacement during exercise should: 1) taste good; 2) not cause gastrointestinal discomfort; 3) be rapidly emptied from the stomach and absorbed in the intestine; 4) provide energy in the form of carbohydrate and a source of electrolytes. It is for this reason that sports drinks are typically available in a variety of flavours (to meet player taste) and contain three main ingredients: water, carbohydrate, and sodium. The



water and carbohydrate provide fluid and energy respectively, while sodium is included to aid water absorption and retention (Maughan & Murray, 2001).

Both carbohydrate and fluid are important. Beverages with a concentrated form of carbohydrate may slow gastric emptying and reduce water absorption. Therefore, it is usually recommended to combine the ingestion of gels, chews, and other solids with water. If drink contents are too concentrated, water may be drawn out of the interstitial fluid and plasma and into the lumen of the small intestine by osmosis, and fluid delivery will be less effective (Gisolfi *et al.*, 1992). Whether this is a problem depends on the goals for hydration and carbohydrate delivery. It must be emphasised here that the addition of sodium and other electrolytes to sports drinks is to increase palatability, maintain thirst (and therefore promote drinking), help prevent hyponatremia, and increase the rate of water uptake. Replacement of the electrolytes lost in sweat during football exercise (~90 minutes) can normally wait until the post-exercise recovery period, unless the player is a particularly heavy and salty sweater and/or the exercise duration is prolonged and in warm conditions (Baker *et al.*, 2016).

Rehydration after training and matches

Rehydration is an important part of the post-exercise recovery process. In the post-exercise occasion, the ingestion of carbohydrate, fluid, and electrolytes is required for replenishing depleted glycogen stores and rehydration respectively (Shirreffs *et al.*, 1996). If players have accrued a body mass deficit (i.e. they are hypohydrated), they should aim to completely replace fluid and electrolyte losses prior to the start of the next training session or match. In most cases, players can just drink according to thirst, as there is plenty of time to restore fluid balance before the next training session.

If hypohydration is severe (>3-5% of BM) or rapid rehydration is needed (e.g. <24 hours before next training or match or pre-season double training sessions), the replacement of fluid and electrolytes in the post-exercise period takes on greater importance. The main factors influencing the effectiveness of post-exercise rehydration are the volume and composition of the fluid consumed.

The recommendation is to drink ~1.5 L of fluid for each 1 kg of body mass deficit (Shirreffs & Sawka, 2011). This is because some of the ingested fluid will be excreted in urine, and studies indicate that ingestion of 150% or more of weight loss is required to achieve normal hydration within 6 hours following exercise (Shirreffs & Maughan, 2000). In most other situations, water and sodium can be consumed with normal eating and drinking practices with no urgency.

Drinking plain water is not the ideal post-exercise rehydration beverage when rapid and complete restoration of body fluid balance is necessary and where all intake is in liquid form (Snell *et al.*, 2010).



Ingestion of water alone in the post-exercise period results in a rapid fall of the plasma sodium concentration and the plasma osmolality. These changes have the effect of reducing the stimulation to drink and increasing the urine output, both of which will delay the rehydration process. Plasma volume is more rapidly and completely restored in the post-exercise period if sodium chloride (77 mmol/L or 450 mg/L) is added to the water consumed (Maughan & Leiper, 1995; Snell *et al.*, 2010). This sodium concentration is similar to the upper limit of the sodium concentration found in sweat but is considerably higher than the sodium concentration of many commercially available sports drinks, which usually contain 10-25 mmol/L (60-150 mg/L). Thus, drinking a beverage containing sodium or eating sodium-containing foods with fluids is recommended during the post-exercise occasion (Maughan *et al.*, 1996; Shirreffs & Sawka, 2011).

The inclusion of potassium in the beverage consumed after exercise would be expected to enhance the replacement of intracellular water and thus promote rehydration. However, there is little experimental evidence to support this. The rehydration drink should also contain carbohydrate (glucose or glucose polymers) because the presence of glucose will also stimulate fluid absorption in the gut and improve beverage taste. Following exercise, the uptake of glucose into the muscle for glycogen re-synthesis should also promote intracellular rehydration (Gisolfi *et al.*, 2001; Gisolfi *et al.*, 1992). Therefore, fluid in the post-exercise occasion can be combined with the ingestion of protein and carbohydrate, which are also required for the player's recovery and adaptation (Hobson & James, 2015; Ivy, 2001; James *et al.*, 2011; James *et al.*, 2013).

Table 2: Fluid guidelines

Occasion	Targets	Principle
Daily intake	Adjust to daily levels of physical activity	Fluid can be ingested via beverages, fruits and vegetable
Prior to exercise	5-7 ml/kg BM	Drink 3-4 before exercise
During exercise	Routine ingestion of fluid	Base intake on individual fluid losses and adjust for exercise intensity and environmental conditions
Post exercise	If rapid rehydration is required, drink 150% of BM losses in the 5 h after the match Avoid alcohol (no more than 1 unit)	Replace fluid losses as a consequence of sweating during exercise. The addition of sodium to the recovery beverage will help with fluid retention and maintain the drive to drink

Source: own source.

Fluid recommendations should be refined with individual sweat rates, specific training intensities, and environmental conditions. Feedback from training/competition performance is required. Fluid needs will increase during training and matches in warmer environments.

ALCOHOL



Although less common in professional clubs, during the season, players may choose to drink alcohol in social settings with teammates, friends and family. There are many cultural or societal factors that may influence this choice, as well as players selecting alcoholic beverages as a means to relieve stress following a match. Moderate/occasional ingestion of alcohol in small quantities (less than 2 units/day) may not be harmful. Nonetheless, alcohol ingestion during the recovery from training and matches can impair liver and glycogen resynthesis, myofibrillar protein synthesis, and rehydration (Collins *et al.* 2021). Furthermore, excessive alcohol intake is also likely to impair physical performance the following day. Therefore, it is generally advised that players should avoid alcohol intake during key periods of the season and following matches when recovery is a priority. There are plenty of social beverage options available, as well as non-alcoholic alternatives, which the sports nutritionist can recommend if players are seeking options.

Summary

- Players should aim to commence training and matches euhydrated.
- Player's fluid loss during training and matches can be monitored and recorded.
- There is a large range in sweat rates among individual players, even in response to the same training session or match.
- Fluid intake should be modified depending on the individual player, the intensity of exercise, and the environmental conditions.
- Players should be educated on fluid needs before, during, and after training and matches



REFERENCES

- Academy of Nutrition and Dietetics, Dietitians of Canada, & American College of Sports Medicine (ACSM).** (2016). Position of the Academy of Nutrition and Dietetics, Dietitians of Canada, and the American College of Sports Medicine: Nutrition and Athletic Performance. *Canadian Journal of Dietetic Practice and Research*, 77(1), 54. [10.1016/j.jand.2015.12.006](https://doi.org/10.1016/j.jand.2015.12.006).
- Aldous, J. W., Christmas, B. C., Akubat, I., Dascombe, B., Abt, G., & Taylor, L.** (2015). Hot and Hypoxic Environments Inhibit Simulated Soccer Performance and Exacerbate Performance Decrements When Combined. *Frontiers in Physiology*, 6, 421. [10.3389/fphys.2015.00421](https://doi.org/10.3389/fphys.2015.00421)
- Ali, A., & Williams, C.** (2013). Isokinetic and isometric muscle function of the knee extensors and flexors during simulated soccer activity: effect of exercise and dehydration. *Journal of Sports Sciences*, 31(8), 907–916. doi:10.1080/02640414.2012.753635
- Aragón-Vargas, L. F., Moncada-Jiménez, J., Hernández-Elizondo, J., Barrenechea, C. A., & Monge-Alvarado, A.** (2009). Evaluation of pre-game hydration status, heat stress, and fluid balance during professional soccer competition in the heat. *European Journal of Sport Science*, 9(5), 269–276. [10.1080/17461390902829242](https://doi.org/10.1080/17461390902829242)
- Armstrong, L. E., Costill, D. L., & Fink, W. J.** (1985). Influence of diuretic-induced dehydration on competitive running performance. *Medicine & Science in Sports & Exercise*, 17(4), 456–461. [10.1249/00005768-198508000-00009](https://doi.org/10.1249/00005768-198508000-00009)
- Armstrong, L. E., Maresh, C. M., Gabaree, C. V., Hoffman, J. R., Kavouras, S. A., Kenefick, R. W., & Ahlquist, L. E.** (1997). Thermal and circulatory responses during exercise: effects of hypohydration, dehydration, and water intake. *Journal of Applied Physiology* (1985), 82(6), 2028–2035. [10.1152/jappl.1997.82.6.2028](https://doi.org/10.1152/jappl.1997.82.6.2028).
- Armstrong, L. E., Pumerantz, A. C., Fiala, K. A., Roti, M. W., Kavouras, S. A., Casa, D. J., & Maresh, C. M.** (2010). Human hydration indices: acute and longitudinal reference values. *International Journal of Sport Nutrition and Exercise Metabolism*, 20(2), 145–153. [10.1123/ijsnem.20.2.145](https://doi.org/10.1123/ijsnem.20.2.145).
- Armstrong, L. E., Soto, J. A., Hacker, F. T., Jr., Casa, D. J., Kavouras, S. A., & Maresh, C. M.** (1998). Urinary indices during dehydration, exercise, and rehydration. *International Journal of Sport Nutrition*, 8(4), 345–355. [10.1123/ijsn.8.4.345](https://doi.org/10.1123/ijsn.8.4.345).
- Badham, L., Stern, S. E., O'Connor, F. K., Wijekulasuriya, G. A., Corcoran, G., Cox, G. R., & Coffey, V. G.** (2023) Fluid intake is a strong predictor of outdoor team sport pre-season training performance. *Journal of Sports Sciences*, 41(1), 1-7. [10.1080/02640414.2023.2191093](https://doi.org/10.1080/02640414.2023.2191093).



- Baker, L. B., & Jeukendrup, A. E.** (2014). Optimal composition of fluid-replacement beverages. *Comprehensive Physiology*, 4(2), 575–620. doi:10.1002/cphy.c130014.
- Baker, L. B., Conroy, D. E., & Kenney, W. L.** (2007). Dehydration impairs vigilance-related attention in male basketball players. *Medicine & Science in Sports & Exercise*, 39(6), 976–983. doi:10.1080/02640414.2015.1055291. [10.1007/s40279-017-0738-7](https://doi.org/10.1007/s40279-017-0738-7)
- Baker, L. B., Dougherty, K. A., Chow, M., & Kenney, W. L.** (2007). Dehydration impairs vigilance-related attention in male basketball players. *Medicine & Science in Sports & Exercise*, 39(7), 1114–1123. doi:10.1097/mss.0b013e3180471ff2
- Baker, L. B., Dougherty, K. A., Chow, M., & Kenney, W. L.** (2007). Progressive dehydration causes a progressive decline in basketball skill performance. *Medicine & Science in Sports & Exercise*, 39(7), 1114–1123. doi:10.1249/mss.0b013e3180574b02
- Baker, L. B., & Jeukendrup, A. E.** (2014). Optimal composition of fluid-replacement beverages. *Comprehensive Physiology*, 4(2), 575–620. doi:10.1002/cphy.c130014
- Baker, L. B., Rollo, I., Stein, K. W., & Jeukendrup, A. E.** (2015). Acute Effects of Carbohydrate Supplementation on Intermittent Sports Performance. *Nutrients*, 7(7), 5733–5763. doi:10.3390/nu7075249
- Baker, L. B., Stofan, J. R., Hamilton, A. A., & Horswill, C. A.** (2009). Comparison of regional patch collection vs. whole body washdown for measuring sweat sodium and potassium loss during exercise. *Journal of Applied Physiology*, 107(3), 887–895. [10.1152/jappphysiol.00197.2009](https://doi.org/10.1152/jappphysiol.00197.2009)
- Baker, L. B., Ungaro, C. T., Barnes, K. A., Nuccio, R. P., Reimel, A. J., & Stofan, J. R.** (2014). Validity and reliability of a field technique for sweat Na⁺ and K⁺ analysis during exercise in a hot-humid environment. *Physiological Reports*, 2(5), e12007. doi:10.14814/phy2.12007
- Baker, L. B., Model, J. B., Barnes, K. A., Anderson, M. L., Lee, S. P., Lee, K. A., Brown, S. D., Reimel, A. J., Roberts, T. J., Nuccio, R. P., Bonsignore, J. L., Ungaro, C. T., Carter, J. M., Li, W., Seib, M. S., Reeder, J. T., Aranyosi, A. J., Rogers, J. A., & Ghaffari, R.** (2020). Skin-interfaced microfluidic system with personalized sweating rate and sweat chloride analytics for sports science applications. *Science Advances*, 6(50), eabe3929. <https://doi.org/10.1126/sciadv.abe3929>
- Below, P. R., Mora-Rodriguez, R., Gonzalez-Alonso, J., & Coyle, E. F.** (1995). Fluid and carbohydrate ingestion independently improve performance during 1 h of intense exercise. *Medicine & Science in Sports & Exercise*, 27(2), 200–210.



- Bendiksen, M., Bischoff, R., Randers, M. B., Mohr, M., Rollo, I., Suetta, C., & Krstrup, P.** (2012). The Copenhagen Soccer Test: physiological response and fatigue development. *Medicine & Science in Sports & Exercise*, *44*(8), 15951603. doi:10.1249/MSS.0b013e31824cc23b
- Cheung, S. S.** (2010). Interconnections between thermal perception and exercise capacity in the heat. *Scandinavian Journal of Medicine & Science in Sports*, *20*(Suppl. 3), 53-59. doi:10.1111/j.1600-0838.2010.01209.x
- Cheuvront, S. N., Carter, R., 3rd, Castellani, J. W., & Sawka, M. N.** (2005). Hypohydration impairs endurance exercise performance in temperate but not cold air. *Journal of Applied Physiology* (1985), *99*(5), 1972-1976. doi:10.1152/jappphysiol.00329.2005
- Cheuvront, S. N., Fraser, C. G., Kenefick, R. W., Ely, B. R., & Sawka, M. N.** (2011). Reference change values for monitoring dehydration *Clinical Chemistry and Laboratory Medicine*, *49*(6), 1033-1037. doi:10.1515/CCLM.2011.170
- Cheuvront, S. N., Haymes, E. M., & Sawka, M. N.** (2002). Comparison of sweat loss estimates for women during prolonged high-intensity running. *Medicine & Science in Sports & Exercise*, *34*(8), 1344-1350. [10.1097/00005768-200208000-00017](https://doi.org/10.1097/00005768-200208000-00017)
- Cheuvront, S. N., & Kenefick, R. W.** (2014). Dehydration: physiology, assessment, and performance effects. *Comprehensive Physiology*, *4*(1), 257-285. doi:10.1002/cphy.c130017
- Cheuvront, S. N., Kenefick, R. W., Charkoudian, N., & Sawka, M. N.** (2013). Physiologic basis for understanding quantitative dehydration assessment. *Comprehensive Physiology*, *4*(1), 455-462. doi:10.3945/ajcn.112.044172
- Cheuvront, S. N., & Montain, S. J.** (2017). Myths and methodologies: Making sense of exercise mass and water balance. *Experimental Physiology*, *102*(9), 1047-1053. doi:10.1113/EP086284
- Collins, J., Maughan, R. J., Gleeson, M., Bilborough, J., Jeukendrup, A., Morton, J. P., Phillips, S. M., Armstrong, L., Burke, L. M., Close, G. L., Duffield, R., Larson-Meyer, E., Louis, J., Medina, D., Meyer, F., Rollo, I., Sundgot-Borgen, J., Wall, B. T., Boulosa, B., Dupont, G., Lizarraga, A., Res, P., Bizzini, M., Castagna, C., Cowie, C. M., D'Hooghe, M., Geyer, H., Meyer, T., Papadimitriou, N., Vouillamoz, M., & McCall, A.** (2021). UEFA expert group statement on nutrition in elite football. Current evidence to inform practical recommendations and guide future research. *British Journal of Sports Medicine*, *55*(8), 416. [10.1097/00005768-200208000-00017](https://doi.org/10.1097/00005768-200208000-00017)
- Coyle, E. F., & Gonzalez-Alonso, J.** (2001). Cardiovascular drift during prolonged exercise: new perspectives. *Exercise and Sport Sciences Reviews*, *29*(2), 88-92. [10.1097/00003677-200104000-00009](https://doi.org/10.1097/00003677-200104000-00009)



- Duffield, R., McCall, A., Coutts, A. J., & Peiffer, J. J.** (2012). Hydration, sweat and thermoregulatory responses to professional football training in the heat. *Journal of Sports Sciences*, 30(10), 957-965. doi:10.1080/02640414.2012.689432
- Eklblom, B.** (1986). Applied physiology of soccer. *Sports Medicine*, 3(1), 50-60. [10.2165/00007256-198603010-00005](https://doi.org/10.2165/00007256-198603010-00005)
- Fleming, J., & James, L. J.** (2014). Repeated familiarisation with hypohydration attenuates the performance decrement caused by hypohydration during treadmill running. *Applied Physiology, Nutrition, and Metabolism*, 39(2), 124-129. doi:10.1139/apnm-2013-0044
- Ganio, M. S., Armstrong, L. E., Casa, D. J., McDermott, B. P., Lee, E. C., Yamamoto, L. M., Marzano, S., Lopez, R. M., Jimenez, L., Le Belleo, L., Chevillotte, E., & Lieberman, H. R.** (2011). Mild dehydration impairs cognitive performance and mood of men. *The British Journal of Nutrition*, 106(10), 1535-1543. doi:10.1017/S0007114511002005
- Gisolfi, C. V., Lambert, G. P., & Summers, R. W.** (2001). Intestinal fluid absorption during exercise: role of sport drink osmolality and [Na⁺]. *Medicine & Science in Sports & Exercise*, 33(6), 907-915. [10.1097/00005768-200106000-00009](https://doi.org/10.1097/00005768-200106000-00009).
- Gisolfi, C. V., Summers, R. W., Schedl, H. P., & Bleiler, T. L.** (1992). Intestinal water absorption from select carbohydrate solutions in humans. *Journal of Applied Physiology*, 73(5), 2142-2150. [10.1152/jappl.1992.73.5.2142](https://doi.org/10.1152/jappl.1992.73.5.2142).
- Gleeson, M.** (1998). Temperature regulation during exercise. *International Journal of Sports Medicine*, 19(Suppl. 2), S96-S99. doi:10.1055/s-2007-971967
- Gonzalez-Alonso, J., Mora-Rodriguez, R., Below, P. R., & Coyle, E. F.** (1995). Dehydration reduces cardiac output and increases systemic and cutaneous vascular resistance during exercise. *Journal of Applied Physiology*, 79(5), 1487-1496. [10.1152/jappl.1995.79.5.1487](https://doi.org/10.1152/jappl.1995.79.5.1487).
- Goulet, E. D.** (2011). Effect of exercise-induced dehydration on time-trial exercise performance: a meta-analysis. *The British Journal of Sports Medicine*, 45(14), 1149-1156. doi:10.1136/bjism.2010.077966
- Hargreaves, M., Dillo, P., Angus, D., & Febbraio, M.** (1996). Effect of fluid ingestion on muscle metabolism during prolonged exercise. *Journal of Applied Physiology*, 80(1), 363-366. [10.1152/jappl.1996.80.1.363](https://doi.org/10.1152/jappl.1996.80.1.363).
- Harper, L. D., West, D. J., Stevenson, E., & Russell, M.** (2014). Technical performance reduces during the extra-time period of professional soccer match-play. *PLoS One*, 9(10), e110995. doi:10.1371/journal.pone.0110995



- Hawley, J. A., Dennis, S. C., & Noakes, T. D.** (1994). Carbohydrate, fluid, and electrolyte requirements of the soccer player: a review. *International Journal of Sport Nutrition*, 4(3), 221-236. doi:10.1123/ijnsn.4.3.221.
- Hobson, R., & James, L.** (2015). The addition of whey protein to a carbohydrate-electrolyte drink does not influence post-exercise rehydration. *Journal of Sports Sciences*, 33(1), 77-84. doi:10.1080/02640414.2014.925570
- Horowitz, M.** (2014). Heat acclimation, epigenetics, and cytoprotection memory. *Comprehensive Physiology*, 4(1), 199-230. doi:10.1002/cphy.c130025
- Impey, S. G., Hammond, K. M., Shepherd, S. O., Sharples, A. P., Stewart, C., Limb, M., & Morton, J. P.** (2016). Fuel for the work required: a practical approach to amalgamating train-low paradigms for endurance athletes. *Physiological Reports*, 4(10), e12803. doi:10.14814/phy2.12803
- Ivy, J. L.** (2001). Dietary strategies to promote glycogen synthesis after exercise. *Canadian Journal of Applied Physiology*, 26(Suppl.), S236-S245. doi:10.1139/h2001-058.
- Jain A.** (2015) Body fluid composition. *Pediatrics in Review*, 36(4), 141-150; quiz 151-152. doi:10.1542/pir.36-4-141. PMID: 25834218.
- James, L. J., Clayton, D., & Evans, G. H.** (2011). Effect of milk protein addition to a carbohydrate-electrolyte rehydration solution ingested after exercise in the heat. *The British Journal of Nutrition*, 105(3), 393-399. doi:10.1017/S0007114510003545
- James, L. J., Evans, G. H., Madin, J., Scott, D., Stepney, M., Harris, R., Stone, R., & Clayton, D. J.** (2013). Effect of varying the concentrations of carbohydrate and milk protein in rehydration solutions ingested after exercise in the heat. *The British Journal of Nutrition*, 110(7), 1285-1291. doi:10.1017/S0007114513000536
- James, L. J., Funnell, M. P., James, R. M., & Mears, S. A.** (2019). Does Hypohydration Really Impair Endurance Performance? Methodological Considerations for Interpreting Hydration Research. *Sports Medicine*, 49(Suppl. 2), 103-114. doi:10.1007/s40279-019-01188-5.
- Jeukendrup, A.** (2014). A step towards personalized sports nutrition: carbohydrate intake during exercise. *Sports Medicine*, 44(Suppl. 1), S25-33. doi:10.1007/s40279-014-0148-z
- Jeukendrup, A. E.** (2011). Nutrition for endurance sports: marathon, triathlon, and road cycling. *Journal of Sports Sciences*, 29(Suppl. 1), S91-99. doi:10.1080/02640414.2011.610348
- Kavouras, S. A., Arnaoutis, G., Makrillos, M., Garagouni, C., Nikolaou, E., Chira, O., & Sidossis, L. S.** (2012). Educational intervention on water intake improves hydration status and



enhances exercise performance in athletic youth. *Scandinavian Journal of Medicine & Science in Sports*, 22(5), 684-689. doi:10.1111/j.1600-0838.2011.01296.x

Kenefick, R. (2018). Fluid Intake Strategies for Optimal Hydration and Performance: Planned Drinking vs. Drinking to Thirst. *Sports Science Exchange*, 29(182), 1-6.

Kilding, A. E., Tunstall, H., Wraith, E., Good, M., Gammon, C., & Smith, C. (2009). Sweat rate and sweat electrolyte composition in international female soccer players during game specific training. *International Journal of Sports Medicine*, 30(6), 443-447. doi:10.1055/s-0028-1105945

Knechtle, B., Knechtle, P., Wirth, A., Rust, C. A., & Rosemann, T. (2012). A faster running speed is associated with a greater body weight loss in 100-km ultra-marathoners. *Journal of Sports Sciences*, 30(11), 1131-1140. doi:10.1080/02640414.2012.692479

Knechtle, B., Rust, C. A., Knechtle, P., & Rosemann, T. (2012). Does Muscle Mass Affect Running Times in Male Long-distance Master Runners? *Asian Journal of Sports Medicine*, 3(4), 247-256. doi:10.5812/asjms.34547.

Lambert, G. P., Lang, J., Bull, A., Eckerson, J., Lanspa, S., & O'Brien, J. (2008). Fluid tolerance while running: effect of repeated trials. *International Journal of Sports Medicine*, 29(11), 878-882. doi:10.1055/s-2008-1038620

Logan-Sprenger, H. M., Heigenhauser, G. J., Jones, G. L., & Spriet, L. L. (2015). The effect of dehydration on muscle metabolism and time trial performance during prolonged cycling in males. *Physiological Reports*, 3(8). doi:10.14814/phy2.12483

Maughan, R. J., Fenn, C. E., & Leiper, J. B. (1989). Effects of fluid, electrolyte and substrate ingestion on endurance capacity. *European Journal of Applied Physiology and Occupational Physiology*, 58(5), 481-486. doi:10.1007/BF02330701.

Maughan, R. J., & Leiper, J. B. (1995). Sodium intake and post-exercise rehydration in man. *European Journal of Applied Physiology and Occupational Physiology*, 71(4), 311-319. doi:10.1007/BF00240410.

Maughan, R. J., Leiper, J. B., & Shirreffs, S. M. (1996). Restoration of fluid balance after exercise-induced dehydration: effects of food and fluid intake. *Journal of Applied Physiology and Occupational Physiology*, 73(3-4), 317-325. doi:10.1007/BF02425493.

Maughan, R. J., & Murray, R. (2001). *Nutrition in Exercise and Sports Sciences: Sports drinks. Basic sciences and practical aspects*. CRC Press.

Maughan, R. J., & Shirreffs, S. M. (2007a). Nutrition and hydration concerns of the female football player. *The British Journal of Sports Medicine*, 41(Suppl. 1), i60-63. doi:10.1136/bjms.2007.036475



- Maughan, R. J., & Shirreffs, S. M.** (2007b). Nutrition for soccer players. *Current Sports Medicine Reports*, 6(5), 279-280.
- Maughan, R. J., Shirreffs, S. M., & Leiper, J. B.** (2007). Errors in the estimation of hydration status from changes in body mass. *Journal of Sports Sciences*, 25(7), 797-804. doi:10.1080/02640410600875143
- Maughan, R. J., Shirreffs, S. M., Merson, S. J., & Horswill, C. A.** (2005). Fluid and electrolyte balance in elite male football (soccer) players training in a cool environment. *Journal of Sports Sciences*, 23(1), 73-79. doi:10.1080/02640410410001730115.
- Maughan, R. J., Watson, P., Evans, G. H., Broad, N., & Shirreffs, S. M.** (2007). Water balance and salt losses in competitive football. *International Journal of Sport Nutrition and Exercise Metabolism*, 17(6), 583-594. doi:10.1123/ijsnem.17.6.583.
- McConell, G. K., Burge, C. M., Skinner, S. L., & Hargreaves, M.** (1997). Influence of ingested fluid volume on physiological responses during prolonged exercise. *Acta Physiologica Scandinavica*, 160(2), 149-156. doi:10.1046/j.1365-201X.1997.00139.x.
- McGregor, S. J., Nicholas, C. W., Lakomy, H. W., & Williams, C.** (1999). The influence of intermittent high-intensity shuttle running and fluid ingestion on the performance of a football skill. *Journal of Sports Sciences*, 17(11), 895-903. doi:10.1080/026404199365452.
- McKenna, K., & Thompson, C.** (1998). Osmoregulation in clinical disorders of thirst appreciation. *Clinical Endocrinology*, 49(2), 139-152.
- Mohr, M., & Krstrup, P.** (2013). Heat stress impairs repeated jump ability after competitive elite soccer games. *Journal of Strength and Conditioning Research*, 27(3), 683-689. doi:10.1097/JSC.0b013e31825c3266
- Montain, S. J., & Coyle, E. F.** (1992). Influence of graded dehydration on hyperthermia and cardiovascular drift during exercise. *Journal of Applied Physiology*, 73(4), 1340-1350. doi:10.1152/jappl.1992.73.4.1340.
- Morris, J. G., Nevill, M. E., Boobis, L. H., Macdonald, I. A., & Williams, C.** (2005). Muscle metabolism, temperature, and function during prolonged, intermittent, high-intensity running in air temperatures of 33 degrees and 17 degrees C. *International Journal of Sports Medicine*, 26(10), 805-814. doi:10.1055/s-2005-837448
- Murray, R.** (2006). Training the gut for competition. *Current Sports Medicine Reports*, 5(3), 161-164. doi:10.1097/01.csmr.0000306307.10697.77.
- Nicholas, C. W., Williams, C., Lakomy, H. K., Phillips, G., & Nowitz, A.** (1995). Influence of ingesting a carbohydrate-electrolyte solution on endurance capacity during intermittent,



high-intensity shuttle running. *Journal of Sports Sciences*, 13(4), 283-290.
doi:10.1080/02640419508732241.

Nybo, L., Rasmussen, P., & Sawka, M. N. (2014). Performance in the heat-physiological factors of importance for hyperthermia-induced fatigue. *Comprehensive Physiology*, 4(2), 657-689.
doi:10.1002/cphy.c130012

Owen, J. A., Kehoe, S. J., & Oliver, S. J. (2013). Influence of fluid intake on soccer performance in a temperate environment. *Journal of Sports Sciences*, 31(1), 1-10.
doi:10.1080/02640414.2012.720701

Popkin, B. M., D'Anci, K. E., & Rosenberg, I. H. (2010). Water, hydration, and health. *Nutrition Reviews*, 68(8), 439-458. doi:10.1111/j.1753-4887.2010.00304.x

Robinson, T. A., Hawley, J. A., Palmer, G. S., Wilson, G. R., Gray, D. A., Noakes, T. D., & Dennis, S. C. (1995). Water ingestion does not improve 1-h cycling performance in moderate ambient temperatures. *European Journal of Applied Physiology and Occupational Physiology*, 71(2-3), 153-160. doi:10.1007/BF00854973.

Rollo, I. (2014). Carbohydrate: the football fuel. *Sports Science Exchange*, 27(127), 1-8.

Rollo, I., Randell, R. K., Baker, L., Leyes, J. Y., Medina Leal, D., Lizarraga, A., Mesalles, J., Jeukendrup, A. E., James, L. J., & Carter, J. M. (2021). Fluid Balance, Sweat Na⁺ Losses, and Carbohydrate Intake of Elite Male Soccer Players in Response to Low and High Training Intensities in Cool and Hot Environments. *Nutrients*, 13(2), 401.
doi:10.3390/nu13020401

Russell, M., & Kingsley, M. (2014). The efficacy of acute nutritional interventions on soccer skill performance. *Sports Medicine*, 44(7), 957-970. doi:10.1007/s40279-014-0184-8

Rust, C. A., Knechtle, B., Knechtle, P., Wirth, A., & Rosemann, T. (2012). Body mass change and ultraendurance performance: a decrease in body mass is associated with an increased running speed in male 100-km ultramarathoners. *International Journal of Sports Medicine*, 33(10), 1505-1516. doi:10.1519/JSC.0b013e318231a7b5

Savoie, F. A., Kenefick, R. W., Ely, B. R., Chevront, S. N., & Goulet, E. D. (2015). Effect of Hypohydration on Muscle Endurance, Strength, Anaerobic Power and Capacity and Vertical Jumping Ability: A Meta-Analysis. *Sports Medicine*, 45(8), 1207-1227.
doi:10.1007/s40279-015-0349-0

Sawka, M. N., Burke, L. M., Eichner, E. R., Maughan, R. J., Montain, S. J., & Stachenfeld, N. S. (2007). American College of Sports Medicine position stand. Exercise and fluid replacement. *Medicine & Science in Sports & Exercise*, 39(2), 377-390.
doi:10.1249/mss.0b013e31802ca597



- Sawka, M. N., Cheuvront, S. N., & Kenefick, R. W.** (2012). High skin temperature and hypohydration impair aerobic performance. *Experimental Physiology*, *97*(3), 327-332. doi:10.1113/expphysiol.2011.061002
- Sawka, M. N., & Greenleaf, J. E.** (1992). Current concepts concerning thirst, dehydration, and fluid replacement: overview. *Medicine & Science in Sports & Exercise*, *24*(6), 643-644.
- Sawka, M. N., Leon, L. R., Montain, S. J., & Sonna, L. A.** (2011). Integrated physiological mechanisms of exercise performance, adaptation, and maladaptation to heat stress. *Comprehensive Physiology*, *1*(4), 1883-1928. doi:10.1002/cphy.c100082
- Shirreffs, S. M., Aragon-Vargas, L. F., Chamorro, M., Maughan, R. J., Serratos, L., & Zachwieja, J. J.** (2005). The sweating response of elite professional soccer players to training in the heat. *International Journal of Sports Medicine*, *26*(2), 90-95. doi:10.1055/s-2004-821112.
- Shirreffs, S. M., Armstrong, L. E., & Cheuvront, S. N.** (2004). Fluid and electrolyte needs for preparation and recovery from training and competition. *Journal of Sports Sciences*, *22*(1), 57-63. doi:10.1080/0264041031000140572
- Shirreffs, S. M., & Maughan, R. J.** (2000). Rehydration and recovery of fluid balance after exercise. *Exercise and Sport Sciences Reviews*, *28*(1), 27-32.
- Shirreffs, S. M., & Sawka, M. N.** (2011). Fluid and electrolyte needs for training, competition, and recovery. *Journal of Sports Sciences*, *29*(Suppl. 1), S39-46. doi:10.1080/02640414.2011.614269
- Shirreffs, S. M., Taylor, A. J., Leiper, J. B., & Maughan, R. J.** (1996). Post-exercise rehydration in man: effects of volume consumed and drink sodium content. *Medicine & Science in Sports & Exercise*, *28*(10), 1260-1271. doi:10.1097/00005768-199610000-00009.
- Snell, P. G., Ward, R., Kandaswami, C., & Stohs, S. J.** (2010). Comparative effects of selected non-caffeinated rehydration sports drinks on short-term performance following moderate dehydration. *Journal of the International Society of Sports Nutrition*, *7*, 28. doi:10.1186/1550-2783-7-28
- Tarnowski, C. A., Rollo, I., Carter, J. M., Lizarraga-Dallo, M. A., Oliva, M. P., Clifford, T., James, L. J., & Randell, R. K.** (2022) Fluid Balance and Carbohydrate Intake of Elite Female Soccer Players during Training and Competition. *Nutrients*, *14*(15), 3188. doi:10.3390/nu14153188.
- Thomas, D. T., Erdman, K. A., & Burke, L. M.** (2016). Position of the Academy of Nutrition and Dietetics, Dietitians of Canada, and the American College of Sports Medicine: Nutrition and Athletic Performance. *Journal of the Academy of Nutrition and Dietetics*, *116*(3), 501-528. doi:10.1016/j.jand.2015.12.006



- Viitasalo, J. T., Kyrolainen, H., Bosco, C., & Alen, M.** (1987). Effects of rapid weight reduction on force production and vertical jumping height. *International Journal of Sports Medicine*, 8(4), 281-285. doi:10.1055/s-2008-1025670
- Watson, G., Judelson, D. A., Armstrong, L. E., Yeargin, S. W., Casa, D. J., & Maresh, C. M.** (2005). Influence of diuretic-induced dehydration on competitive sprint and power performance. *Medicine & Science in Sports & Exercise*, 37(7), 1168-1174. doi:10.1249/01.mss.0000170100.93279.be.
- Wenger, C. B.** (1972). Heat of evaporation of sweat: thermodynamic considerations. *Journal of Applied Physiology*, 32(4), 456-459. doi:10.1152/jappl.1972.32.4.456
- Williams, C., & Rollo, I.** (2015). Carbohydrate Nutrition and Team Sport Performance. *Sports Medicine*, 45(Suppl. 1), S13-22. doi:10.1007/s40279-015-0399-3

